

AD-A118 689

ALBERTA UNIV EDMONTON DEPT OF CHEMISTRY

F/G 7/4

SIMULATION OF THE DROPPING MERCURY ELECTRODE BY ORTHOGONAL COLL--ETC(U)

AUG 82 S PONS, B SPEISER, J MCALLEER

N00014-82-G-0017

UNCLASSIFIED TR-13

NL

END
DATE
FILED
9 82
DTB

AD A118689

DTIC FILE COPY

OFFICE OF NAVAL RESEARCH
Contract N00014-82-G-0017
Task No. NR 359-718
TECHNICAL REPORT NO. 13

Simulation of the Dropping Mercury Electrode by Orthogonal Collo-
cation.

By

Stanley Pons, Bernd Speiser, J. McAleer, and P.P. Schmidt**

Prepared for Publication in

Electrochimica Acta

University of Alberta
Department of Chemistry
Edmonton, Alberta, Canada
T6G 2G2

**Oakland University
Department of Chemistry
Rochester, Michigan

August 18, 1982

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC
ELECTE
AUG 27 1982
S H D

(2)

REPORT DOCUMENTATION PAGE	
1. REPORT NUMBER	11
2. GOVT ACCESSION NO.	AD-A118689
3. TITLE (and subtitle)	Simulation of the Dropping Mercury Electrode by Orthogonal Collo- cation.
4. AUTHOR(S)	Stanley Pons, Bernd Speiser and J. McAleer
5. PERFORMING ORGANIZATION NAME AND ADDRESS	Department of Chemistry University of Alberta Edmonton, Alberta, Canada T6G 2G2
6. CONTROLLING OFFICE NAME AND ADDRESS	Office of Naval Research Chemistry Program Arlington, Virginia 22217 - Chemistry Code 472
7. DISTRIBUTION STATEMENT (or MR. Report)	This document has been approved for public release and sale; its distribution unlimited.
8. NUMBER OF PAGES	25
9. SECURITY CLASSIFICATION	Unclassified
10. KEY WORDS (Continue on reverse side if necessary and identify by block number)	Electrochemistry, Mercury Electrode, Simulation, Orthogonal Collo- cation.
11. ABSTRACT (Continue on reverse side if necessary and identify by block number)	The theory for the application of orthogonal collocation techniques to the dropping mercury electrode is given. The technique is shown to adapt readily to non-linear geometries.
12. AD INSTRUCTIONS REFINERY CODES, ETC. (check)	
13. HELPLINE & CATALOG NUMBER	
14. TYPE OF REPORT & PERIOD COVERED	Technical Report - 13
15. PERFORMING ORG. REPORT NUMBER	N00014-82-G-0017
16. CONTRACT OR GRANT NUMBER(S)	
17. ORIGINAL EDITION NUMBER, TASK AREA & WORK UNIT NUMBER	Task No. NR 359-718
18. REPORT DATE	August 18, 1982
19. NUMBER OF PAGES	25
20. SECURITY CLASSIFICATION/DECLASSIFICATION RESCALING	

DD FORM 1473 EDITION OF 10-65 10-65
5-74 6102 11 814 2401 Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (If other than Report)

82 8 27 051

Orthogonal collocation techniques are used to simulate the solution to the hydrodynamic partial differential equations describing mass transport to a dropping mercury electrode (DME). Accurate values for the concentration profiles and current are obtained with minimal computational effort.

Simulation of the Dropping Mercury Electrode
by Orthogonal Collocation

Stanley Pons¹, Bernd Speiser¹, Jerome P. McAleer¹
and P.P. Schmidt²

¹Department of Chemistry
University of Alberta
Edmonton, Alberta, Canada
T6G 2G2

and

²Department of Chemistry
Oakland University
Rochester, Michigan 48063

^{*}To whom proofs should be sent.



14

The weighted residual method of orthogonal collocation has been used to generate accurate solutions to electrochemical diffusion problems (1-11). The purpose of this work is to demonstrate the utility of the method by application directly to a differential equation containing convective terms. The case considered here is the diffusive and convective transport of an electroactive species to the dropping mercury electrode (DME).

In terms of radial distance r from the center of the mercury drop, the hydrodynamic equation describing the concentration of the electroactive species in time is given (12) by

$$\frac{\partial c}{\partial t} = D \frac{\partial^2 c}{\partial r^2} + 2D \frac{\partial c}{\partial r} - v_r \frac{\partial c}{\partial r} \quad (1)$$

where v_r is the velocity of a point in solution toward the growing mercury drop. The radius r_0 of the drop grows as

$$r_0 = \left(\frac{3m}{4\pi d} \right)^{1/3} \quad (2)$$

where m is the flow rate and d the density of mercury. Considering only convection, a point r in solution at $t=0$ is pushed outwards. As volume must be conserved, its movement is described by

$$r = \left(\frac{3m(t+t')}{4\pi d} \right)^{1/3} \quad (3)$$

where t' is the time it would take for a sphere of radius r' to be formed at a flow rate m .

The velocity of this point is thus given by

$$v_r = \frac{dr}{dt} = \frac{\left(\frac{3m}{4\pi d} \right)^{1/3}}{3 \left(\frac{3m(t+t')}{4\pi d} \right)^{2/3}} = \frac{3m}{4\pi d} \quad (4)$$

We wish to normalize the distance coordinate to a $[0,1]$ space where 0 represents the electrode surface, and 1 represents a point in solution where no diffusional processes occur during the experiment. We denote this real point as M . Its movement is given by an equation analogous to (3):

$$M = \left(\frac{3m(t+t')}{4\pi d} \right)^{1/3} \quad (5)$$

where t' is the time needed to grow a sphere of radius M . The space between the electrode surface and M then is used in normalization.

It is seen that such a space is time dependent, i.e. the space will diminish in time. This is physically observed as the decrease in diffusion layer thickness resulting in an increase in current at constant potential. For our analysis, it is desired to eliminate this time dependency of the distance coordinate. This is done by observing that the difference in the cubes of the quantities M and r_0 is time independent.

$$M^3 - r_0^3 = \frac{3mt^*}{4\pi d} \quad (6)$$

A point in solution will move with respect to the growing mercury drop surface as well. We can refer to a point fixed in the M^3 -

ORIC Copy Requested	Accession No.
	NTIS Code
	DTIC File Number
	Classification
	Justification
	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
Dist. Avail. and/or Special	Availability Codes
Distr.	By
Distribution/ Availability	

r_0 space as R if R is defined as

$$R = \frac{r - r_0}{(M^3 - r_0^3)^{1/3}} \quad (7)$$

where r is the radial distance from the center of the drop to the point in question.

We define our dimensionless time variable T as

$$T = \frac{tD}{(M^3 - r_0^3)^{2/3}} \quad (8)$$

and a constant s as

$$s = \frac{a^3}{D(M^3 - r_0^3)^{1/3}} \quad (9)$$

where a is given by

$$a = \left(\frac{3M}{4\pi D}\right)^{1/3} \quad (10)$$

We need now to perform the transformation of the differential equation (1) to dimensionless coordinates. In terms of the new coordinates and constants, we find that

$$t = \frac{a^6 T}{D^3 s^2} \quad (11)$$

and

$$r = \frac{Ra^3}{SD} + \frac{a^3 T^{1/3}}{s^{2/3} D} \quad (12)$$

3

4

For a function $w = f(x,y)$ which has continuous first and second partial derivatives, we wish to be able to transform to the new variables u and v where u and v are given by

$$x = \phi(u,v) \quad (13)$$

and

$$y = \psi(u,v) \quad (14)$$

The resulting first partial derivatives are given by

$$\frac{\partial w}{\partial u} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial u} \quad (15)$$

$$\frac{\partial w}{\partial v} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial v} \quad (16)$$

If $w(x,y)$ corresponds to $c(x,t)$, and ψ corresponds to $\frac{a^6 T}{s^2 D^3}$

ψ corresponds to $\frac{a^3 R}{SD} + \frac{a^3 T^{1/3}}{s^{2/3} D}$, then we have

$$\frac{\partial c}{\partial R} = \frac{\partial c}{\partial r} \frac{\partial r}{\partial R} \left(\frac{a^3 R}{SD} + \frac{a^3 T^{1/3}}{s^{2/3} D} \right) + \frac{\partial c}{\partial t} \frac{\partial t}{\partial R} \left(\frac{a^6 T}{D^3 s^2} \right) \quad (17)$$

and

$$\frac{\partial c}{\partial T} = \frac{\partial c}{\partial r} \frac{\partial r}{\partial T} \left(\frac{a^3 R}{SD} + \frac{a^3 T^{1/3}}{s^{2/3} D} \right) + \frac{\partial c}{\partial t} \frac{\partial t}{\partial T} \left(\frac{a^6 T}{D^3 s^2} \right) \quad (18)$$

Thus

$$\frac{\delta c}{\delta t} = \frac{a^3}{8D} \frac{\delta c}{\delta r} ; \quad \frac{\delta c}{\delta r} = \frac{8D}{a^3} \frac{\delta c}{\delta R} \quad (19)$$

$$\frac{\delta c}{\delta t} = \frac{a^3}{3a^{2/3}D^{2/3}} \frac{\delta c}{\delta t} + \frac{a^6}{a^2D^3} \frac{\delta c}{\delta t} ; \quad \frac{\delta c}{\delta t} = \frac{a^2D^3}{a^6} \frac{\delta c}{\delta t} = \frac{a^{4/3}D^2}{3a^3T^{2/3}} \frac{\delta c}{\delta t} \quad (20)$$

Combining (19) and (20), we have

$$\frac{\delta c}{\delta t} = \frac{a^2D^3}{a^6} \frac{\delta c}{\delta t} = \frac{a^{7/3}D^3}{3a^6T^{2/3}} \frac{\delta c}{\delta R} \quad (21)$$

The second partial with respect to the distance coordinate is obtained by recognizing that

$$\frac{\delta^2 c}{\delta r^2} = \frac{\delta}{\delta r} \left(\frac{\delta c}{\delta r} \right) = \frac{1}{\delta r} \left(\frac{\delta D}{\delta r} \frac{\delta c}{\delta R} \right) \quad (22)$$

It can be shown that

$$\frac{1}{\delta r} = \frac{8D}{a^3} \frac{1}{\delta R} \quad (23)$$

so that we now have

$$\frac{\delta^2 c}{\delta r^2} = \frac{a^2D^2}{a^6} \frac{\delta^2 c}{\delta R^2} \quad (24)$$

Equations (12), (19), (21), and (24) are substituted into the differential equation to get

$$\frac{a^2D^3}{a^6} \frac{\delta c}{\delta t} = \frac{a^{7/3}D^3}{a^6T^{2/3}} \frac{\delta c}{\delta R} = \frac{a^2D^3}{a^6} \frac{\delta^2 c}{\delta R^2} + \left[\frac{-2D^3}{Ra^3 + \frac{a^3T^{1/3}}{a^2D}} - \frac{a^3}{3(Ra^3 + \frac{a^3T^{1/3}}{a^2D})^2} \right] \frac{8D}{a^3} \frac{\delta c}{\delta R} \quad (25)$$

where

$$c^0 = \frac{c}{c^b} \quad (26)$$

c^b being the bulk concentration of substrate. Equation (25) is readily reduced to

$$\frac{\delta c}{\delta t} = \frac{a^2c^0}{\delta R^2} + \left[\frac{2}{Ra^{1/3}T^{1/3}} + \frac{a^{1/3}}{3T^{2/3}} - \frac{a}{3(Ra^{1/3}T^{1/3})^2} \right] \frac{\delta c}{\delta R} \quad (27)$$

DISCRETIZATION OF THE DIFFERENTIAL EQUATION

Accurate solutions of the differential equation (27) are obtained at the roots of shifted Jacobi polynomials on the $[0,1]$ interval as a function of time by orthogonal collocation (1,3,11). In terms of the collocation coefficients A_{ij} and B_{ij} (functions of the roots of the polynomial only as previously defined) we have that, at the roots R_i of the polynomial,

$$\frac{\delta c}{\delta R} \bigg|_{R_i} = \sum_{j=1}^{N+2} A_{ij} c^0(R_j, t) \quad (28)$$

and

$$\frac{d^2c^*}{dr^2} \Big|_{R_1} = \sum_{j=1}^{N+2} B_{1j} c^*(R_j, T) \quad (29)$$

where N is the order of the polynomial. $N+2$ terms are taken in order to include the electrode surface ($R=0$) and the point M in solution where no diffusion occurs ($R=1$).

We take as an experiment the imposition of a constant potential to the electrode surface such that $c^*(0, T)$ is equal to zero during the growth of the drop. The mechanism is simple single electron transfer. We refer to the concentration of the substrate only.

We thus recognize that

$$c^*(0, T) = 0 \quad (30)$$

and

$$c^*(1, T) = 1 \quad (31)$$

such that

$$\frac{dc^*}{dr} \Big|_{R_1} = A_{1, N+2} + \sum_{j=2}^{N+1} B_{1j} c^*(R_j, T) \quad (32)$$

and

$$\frac{d^2c^*}{dr^2} \Big|_{R_1} = B_{1, N+2} + \sum_{j=2}^{N+1} B_{1j} c^*(R_j, T) \quad (33)$$

These differentials, now referring to explicit points in simulation space, reduce the partial differential equation to the following set of N first order ordinary differential equation in N unknowns which may be solved simultaneously for the $c(R_j, T)$ (1, 3, 11)

$$\frac{dc^*}{dT} \Big|_{R_1} = B_{1, N+2} + \sum_{j=2}^{N+1} B_{1j} c^*(R_j, T) + \left[\frac{2}{R + \delta^{1/3} \tau^{1/3}} + \frac{\delta^{1/3}}{3\tau^{2/3}} - \frac{\delta}{3(R + \delta^{1/3} \tau^{1/3})^2} \right] \left[A_{1, N+2} + \sum_{j=2}^{N+1} A_{1j} c^*(R_j, T) \right] \quad (34)$$

THE SIMULATED CURRENT

The current is given by Ricker's first law:

$$I = nFAD \left(\frac{dc^*}{dr} \right) \Big|_{r=0} \quad (35)$$

where A is the area of the IME

$$A = 4\pi r_0^2 \quad (36)$$

Substitution of (4) and (19) into (35) and simplifying, results, in terms of the dimensionless parameters,

$$I = 4\pi nFc^* \delta^{-1/3} \tau^{2/3} \left(\frac{dc^*}{dr} \right) \Big|_{r=0} \quad (37)$$

The differential is immediately discretized, considering the

boundary conditions $c^*(0, T) = 0$ and $c^*(1, T) = 1$, to

$$i = 4\pi n F c D a^3 \beta^{-1/3} T^{2/3} \left[A_{1,N+2} + \sum_{j=2}^{N+1} A_{jj} c^*(R_j, T) \right] \quad (38)$$

$$= 21.39 n c b_m \beta^{-1/3} T^{2/3} \left[A_{1,N+2} + \sum_{j=2}^{N+1} A_{jj} c^*(R_j, T) \right] \quad (39)$$

The current is calculated in the output subroutine of the program.

RESULTS

The simulated current is compared to the solution offered by Koutecky (14) which is corrected for spherical diffusion

$$i_d = 708 n c b_m^{2/3} D^{1/2} t^{1/6} (1 + 34.7 b^{1/2} t^{1/6} m^{-1/3} + 100 D t^{1/3} m^{-2/3}) \quad (40)$$

Again, equation (11) is used to convert to dimensionless time

$$i = 708 n c b_m^{1/6} \beta^{-1/3} \left(\frac{3}{4\pi D} \right)^{1/3} (1 + 34.7 \left(\frac{3}{4\pi D} \right)^{1/3} T^{1/6} \beta^{-1/3} + 100 \left(\frac{3}{4\pi D} \right)^{2/3} T^{1/3} \beta^{-2/3}) \quad (41)$$

$$= 18.45 n c b_m T^{1/6} \beta^{-1/3} (1 + 0.902 T^{1/6} \beta^{-1/3} + 0.0678 T^{1/3} \beta^{-2/3}) \quad (42)$$

We may also compare the current to the Ilkovic equation

$$i = 708 n b^{1/2} c b_m^{2/3} t^{1/6} \quad (43)$$

which is converted by the same substitutions to

$$i = 708 n c b_m \left(\frac{3}{4\pi D} \right)^{1/3} T^{1/6} \beta^{-1/3} \quad (44)$$

$$= 18.45 n c b_m^{2/3} T^{1/6} \beta^{-1/3} \quad (45)$$

Values for the parameters for a typical real case were chosen and the results are given in Figure 2.

COMPUTATIONAL ASPECTS

Results accurate to 0.4 % of Koutecky's calculated values are obtained with minimal computational effort. Using 6 solution points (collocation points; polynomial roots) 1000 current values and concentration profiles are generated by the program in less than 0.5 seconds using an Amdahl 470-V6 computer. The program, which has been described (4,13), is modular in design. Switching from the problem described herein to simple linear diffusion requires the changing of only 2 program statements. Such minimal changing of the program is retained with all other electrochemical modeling problems.

CONCLUSIONS

Orthogonal collocation may be adapted in a straightforward manner to provide accurate solutions to hydrodynamic problems in electrochemistry. The problem herein discussed describes only

11

the first drop in an actual experiment and useful data may only be obtained from subsequent drops which apparently encounter reproducible concentration perturbations of the reaction layer. These perturbations are due to convective transport initiated by the first drop. We are preparing the simulation treatment for this case, as well as those describing the current obtained when following chemical reactions and adsorption are involved in the overall reaction mechanism.

12

ACKNOWLEDGEMENTS

Part of this work was supported by a grant from the Office of Naval Research, Washington, and the Department of Chemistry, University of Alberta. The authors thank Professor Fraser Birss for his most helpful comments on the mathematics of the problem.

REFERENCES

1. L. Whiting and P. Carr, *J. Electroanal. Chem.* 81, 1 (1977).
2. S. Pons and P.P. Schmidt, *Electrochim. Acta* 25, 987 (1980).
3. S. Pons, *Electrochim. Acta* 26, 601 (1981).
4. S. Pons in A. Bard, Ed., "Electroanalytical Chemistry", in press.
5. S. Pons, *Can. J. Chem.* 59, 1538 (1981).
6. B. Speiser and S. Pons, *Can. J. Chem.*, in press.
7. B. Speiser, S. Pons, and A. Rieker, *Electrochim. Acta*, in press.
8. B. Speiser and A. Rieker, *J. Electroanal. Chem.* 102, 1 (1979).
9. B. Speiser, *J. Electroanal. Chem.* 110, 69 (1980).
10. B. Speiser and S. Pons, *Electrochim. Acta*, in press.
11. A. Bewick, J. Mellor, and S. Pons, *Electrochim. Acta* 25, 931 (1980).
12. J. Heyrovsky and J. Kuta, "Principles of Polarography", Czechoslovak Academy of Sciences, Prague, 1965.
13. J. Villadsen and M. Michelsen, "Solution of Differential Equation Models by Polynomial Approximation", Prentice Hall, Englewood Cliffs, 1978.
14. J. Koutecky and M.V. Stackelberg in P. Zuman and I. Kolthoff, Eds., "Progress in Polarography", Vol. I, Wiley, New York, 1962.




FIGURE LEGENDS

- (1) The geometry of the growing spherical mercury drop.
- (2) Comparison of (a) simulated results with (b) the Koutecky model and (c) the Ilkovic equation. Parameters: $\beta = 0.35$, $c_b = 10^{-5}$ mol/cm³, $n = 2$ mg/s.

ACKNOWLEDGEMENT

We thank the Office of Naval Research, Washington, D.C., for support of part of this work. X

SP472-3/A1

472:GAN:716:enj
78u+2-60d

SP472-3/A11

472:GAN:716:6dc
78u+2-60b

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-M-A-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCarter San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D One Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. S. Water, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6130 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. M. Orsak Port Hueneme, California 93041	1
The Assistant Secretary of the Navy (R&S) Department of the Navy Room #E730, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 311C (Dr. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 3, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Basmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Sealfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 92133	1
Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112			

TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. Paul Delhey Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 4NN United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106	1	Dr. Sam Parsons Department of Chemistry Purdue University West Lafayette, Indiana	1
Dr. D. W. Benson Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91109	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Ambron Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Warfare Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co., Inc. P.O. Box 306 Sunnyvale, California 94088	1	Dr. J. Boehmer Electrochemicals Corporation Attention: Technical Library 2483 Charlevoix Road Mountain View, California 94030	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 2100 N. Research Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48043	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1	Dr. R. Richter Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1

TECHNICAL REPORT DISTRIBUTION LIST, 359No.
Copies

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, NY 11210 1

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, PR 00931 1

Dr. Joseph Gordon II
IBM Corporation
K33/281
5600 Cottle Road
San Jose, CA 95193 1

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, CA 91103 1

DATE
ILMED
.8